

RESEARCH ARTICLE

Tetrodotoxin and fatty acids contents of *Lagocephalus sceleratus* (Gmelin, 1789) collected in Antalya, Turkey, by MS/MS and GC/MS analyses

**Neşe Kırmır¹, Fatih Göker¹, Elif Özgür Özbek², Burak Coban³,
Esra Billur Balcıoğlu⁴, Bayram Öztürk^{4,5}, Kasım Cemal Güven^{5*}**

¹ Department of Pharmacognosy, Faculty of Pharmacy, Anadolu University, 26470, Eskişehir, TURKEY

² Antalya Metropolitan Municipality, Marine Biology Museum, TURKEY

³ Department of Chemistry, Faculty of Arts and Sciences, Bulent Ecevit University, Zonguldak, 67100, TURKEY

⁴ Faculty of Fisheries, Istanbul University, Istanbul, TURKEY

⁵ Turkish Marine Research Foundation (TUDAV), P.O. Box: 10, Beykoz, Istanbul, TURKEY

*Corresponding author: kcguyen@yahoo.com.tr

Abstract

Tetrodotoxin (TTX) and fatty acids contents of five specimens of *Lagocephalus sceleratus* collected from Antalya on the Turkish Mediterranean coast were studied. TTX was determined by LC-MS/MS analysis in intestines, liver, ovary and muscle. Fatty acids were determined by GC/MS analyses. The oil content was not high (6%) but polyunsaturated n-3 and n-6 ratios were high.

Keywords: Tetrodotoxin, TTX, fatty acids, *Lagocephalus sceleratus*, LC-MS/MS, GC/MS.

Received: 10.11.2016, **Accepted:** 15.12.2016

Introduction

Tetrodotoxin (TTX) has been found in various marine organisms (Miyazawa and Noguchi 2001; Wu *et al.* 2005; Hwang *et al.* 2007) and also *Shewanella* red alga (Simidu *et al.* 1990). It was first isolated by Yokoo (1950) and its structure was elucidated by various authors (Yokoo 1950; Tsuda *et al.* 1964; Woodward 1964; Goto *et al.* 1965). It is a heterocyclic guanidine compound, chemically (4*R*,4*aR*,5*R*,6*S*,7*S*,8*S*,8*aR*,10*S*,12*S*)-2-azaniumylidene-4,6,8,12-tetrahydroxy-6-(hydroxymethyl)-2,3,4,4*a*,5,6,7,8-octahydro-1*H*-8*a*,10-methano-5,7-(epoxymet-

hanooxy) quin-azolin-10-olate. There are a number of natural analogues of TTX including, 4-epiTTX; 6-epiTTX; 11-deoxyTTX; 6,11-dideoxyTTX; 8,11-dideoxyTTX; 11-oxo-TTX; 11-norTTX-6,6-diol; 11-norTTX-6(R)-ol; 11-norTTX-6(S)-ol; chiriquitox; TTX-8-O-hemisuccinate; TTX-11-carboxylic acid (Bane *et al.* 2014).

Since the opening of the Suez Canal in 1869 more than 300 aquatic species have migrated from the Red Sea to the Mediterranean Sea (Bentur *et al.* 2008). Golani listed the migrating fish to the Mediterranean Sea including *Lagocephalus spadiceus*, *L. suenzensis*, *L. sceleratus* in (Antalya) Mediterranean Sea was recorded by Akyol *et al.* (Golani 2002; Akyol *et al.* 2005). TTX is a toxin found in various organs, such as muscle, intestine, liver testis, ovary and skin, of pufferfish *L. sceleratus* (Aydin 2011). The origin of TTX in fish is unknown. In contrary to wild fish, it was not found in cultured fish (Kono *et al.* 2008). The mortality rate is very high, thus it is forbidden to catch and sell this fish, although *L. sceleratus* has been consumed as food in Eastern Asia as well as in the Mediterranean countries (Nader *et al.* 2012; Rodriguez *et al.* 2012). TTX producing 36 bacterial strains has been isolated from the pufferfish (Yu *et al.* 2011).

Various methods have been published on the determination of TTX, among which LC-MS/MS technique most sensitive (Chen and Chou 1998; Jen *et al.* 2008; Jang *et al.* 2010; Chulanetra *et al.* 2011; Lin and Hwang 2012; Rodriguez *et al.* 2012; Silva *et al.* 2012). Biological mouse assay has also been used although not sensitive it is used as an assay for many marine toxic compounds (Brillantes *et al.* 2003; Wu *et al.* 2005; Sabrah *et al.* 2006; Noguchi and Arakawa 2008; Saoudi *et al.* 2008; Simon *et al.* 2009; Köşker *et al.* 2015).

The content of TTX analogues and isomers of *Lagocephalus* sp. and *L. sceleratus* from the Mediterranean Sea were determined by LC-MS/MS (Rodriguez *et al.* 2012; Kızılkaya 2014).

Many papers were published on fatty acids contents of fish oils which are important for human health. Especially n-3 fatty acids reduce the risk of heart attacks (Daviglius *et al.* 1997). Polyunsaturated fatty acids (PUFA) Eicosapentaenoic acid (EPA) 20:5 n-3 and docosohexaenoic acid (DHA) 22:6 n-3: important fatty acids of *L. sceleratus* were investigated in the Mediterranean Sea samples (Özoğul *et al.* 2009; Nurullahoğlu and Ulusoy 2011; Köşker 2014) and 26 species of pufferfish including *Lagosephalus* spp. (*L. inermis*, *L. wheeleri*, *L. glovevi*, *L. guntheri*, *L. spadiceus* and *L. lunaris*) imported from Korea, China and Bangladesh were investigated (Oyaizu *et al.* 2000).

In this paper Tetrodotoxin and fatty acids contents of *Lagocephalus sceleratus* var. in Turkish coast of Mediterranean Sea are reported.

Materials and Methods

Five samples of *L. scleratus* were caught by gillnets and trammel nets in the Gulf of Antalya in February 2014. All captured fishes were transferred to the laboratory maintaining cold chain.

Extraction of TTX from organs and LC-MS/MS conditions for TTX determination

Following the method by Jang *et al.* (2010), 2.6 g ovary, 10 g intestine, 35 g muscle, 1 g liver tissues were extracted. Each sample was homogenized with 0.05 M acetic acid contained methanol in homogenizer and then heated 5 min in water bath, centrifuged at 4000 rpm for 15 min and supernatant phase was separated and filtered. 2 mL aliquot was applied on the reverse phase column C18 (purchased from Macherey-Nagel) which was equilibrated with water after methanol. The first passing solution was discarded and following solution was collected with chloroform. Aqueous phase was separated and applied for LC-MS/MS analysis.

Spectroscopic analysis of aqueous part was performed Applied Biosystems 3200 Q-Trap LC-MS/MS instrument equipped with an ESI ion source with direct-injection to mass spectrometry at positive ionization mode. Enhanced Product Ionization Mode (EPI) chosen for TTX analysis which enabled to select and break a specific molecular weight.

Extraction of fatty acids from flesh

After skin and internal organs were separated, muscle fillets were cut into small pieces using a knife then homogenized using a homogenizer. Homogenized clean spineless fillets (100 g) were extracted with n-hexane using a Soxhlet apparatus for 6 h. The lipid content was determined after removal of solvent and kept at -18 °C. Methyl esters of fatty acids were prepared through modified method of Joseph and Achman (1992) and Aslan *et al.* (2009), and analyzed by GC-MS (Agilent 5975 GC-MSD). Innowax FSC column (60 m x 0.25 mm, 0.25 µm film thickness) was used with helium as carrier gas (0.8 mL min⁻¹). GC oven temperature was kept at 60 °C for 10 min and programmed to 220 °C at a rate of 4 °C min⁻¹ and kept constant at 220 °C for 10 min and then programmed to 240 °C at a rate of 1 °C min⁻¹. Split ratio was adjusted 40:1. The injector temperature was at 250 °C. Ionization energy was 70 eV. Mass range was from m/z 35 to 450. FID temperature was 300 °C.

The amount of each fatty acid was estimated by comparison of their mass spectra with those in the Baser Library of Essential Oil Constituents, Adams Library (Adams 2007), MassFinder Library (Hochmuth 2008), Wiley GC/MS Library (McLafferty and Stauffer 1989) and confirmed by comparison of their retention indices. Alkanes were used as reference points in the calculation of relative retention indices (RRI) (Curvers *et al.* 1985). Relative percentage amounts of the separated compounds were calculated from FID chromatograms.

Results and Discussion

The mass spectra of *L. sceleratus* organs are shown in Figure 1. Various authors reported the similar results on these as 230, 304, 302, 290 (Shoji *et al.* 2001), 320, 302, 251 (Wu *et al.* 2005), 320, 302, 280, 260 (Jen *et al.* 2008), 320, 304, 302, 288 (Jang *et al.* 2010), 320, 302, 284, 256 (Rodriguez *et al.* 2012), 320, 302 (Silva *et al.* 2012), 320, 302 (Chulanetra *et al.* 2011).




Figure 1a. MS spectrum of compound m/z 320 (M+H) extracted from muscle.




Figure 1b. MS spectrum of compound m/z 320 (M+H) extracted from intestine.




Figure 1c. MS spectrum of compound m/z 320 (M+H) extracted from liver.




Figure 1d. MS spectrum of compound m/z 320 (M+H) extracted from ovary.

In summary, the dominant peaks are 320, 302, 290 and 260. The dominant peaks of tetradoxin in examined tissues were found as 320, 302, 280 in muscles, 320, 302, 284, 2 in intestines, 320, 302, 280 in livers, 320, 302, 284, 200 in ovary. The comparison of LC-MS/MS results with the findings of indicated literature proved that muscle, liver, intestine and ovary contain TTX.

Table 1. Fatty acid compositions of flesh (g/100 g lipids)

Fatty acids	g/100 g lipids
C12:0	-
C13:0	-
C14:0	-
C15:0	NSA
C16:0	6.1
C17:0	-
C18:0	6.8
C19:0	-
C20:0	-
C21:0	-
C24:0	-
Σ SFA	12.9
C16:1 (<i>n</i> -7)	NSA
C17:1 (<i>n</i> -7)	-
C18:1 (<i>n</i> -9)	13.7
C18:1 (<i>n</i> -7)	3.2
C20:1 (<i>n</i> -9)	-
C22:1 (<i>n</i> -11)	NSA
C24:1(<i>n</i> -9)	-
Σ MUFA	16.9
C18:2 (<i>n</i> -6)	1.5
C18:3 (<i>n</i> -6)	-
C18:3 (<i>n</i> -3)	-
C18:4 (<i>n</i> -3)	-
C20:2 (<i>n</i> -6)	-
C20:4 (<i>n</i> -6)	6.7
C20:5 (<i>n</i> -3) (EPA)	NSA
C22:5 (<i>n</i> -3)	-
C22:6 (<i>n</i> -3) (DHA)	46.8
Σ PUFA	55.0
Σ Unsat.	71.9
Sat./Unsat.	0.1
Σ MUFA/Σ PUFA	0.31
Σ <i>n</i> -3 PUFA	46.8
Σ <i>n</i> -6 PUFA	4.7
<i>n</i> -3/ <i>n</i> -6	0.9
DHA/EPA	-
Lipid *	0.6
Unidentified	15.2

-: not detected; NSA: no significant amount
(<0.01 g/100 g lipids), * (g/100 g sample)

Fatty acid compositions are shown in Table 1. The lipid level of this species is relatively very low compared to other fish (Love 1970) as 0.6% in our findings, comparable to the other studies reported between 0.2-1.8% (Özoğul *et al.* 2009; Nurullahoğlu and Ulusoy 2011; Aydın *et al.* 2013; Köşker 2014). The recent studies and our result indicate the composition of fatty acids in *L. sceleratus*

collected from Turkish coast of Mediterranean Sea are mainly C16, C18, C18:1 (n-9), C18:1 (n-7), C18:2 (n-6), C20:4 (n-6), C20:5 (n-3) (EPA), C22:5 (n-3), C22:6 (n-3) (DHA), however, the amounts temporally vary as indicated in two reports (Nurulloğlu and Ulusoy 2011; Aydın *et al.* 2013). The total saturated fatty acid (SFA), mono unsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA) were determined as 12.9, 16.9, 55.0%, respectively. Total percentage of omega-3 PUFAs was found as (only DHA was found) 46.8%. Notably C20:5 (n-3) (EPA) was not found significant amount for the recent study unlike to the previous reports. These ratios were comparable with other studies (Özoğul *et al.* 2009; Nurulloğlu and Ulusoy 2011; Aydın *et al.* 2013; Köşker 2014).

Conclusion

TTX was detected in edible muscle, liver, intestine and ovary of *L. sceleratus*. Oil and its fatty acids were determined. It was found that the total lipid content of *L. sceleratus* was low as indicated previously.

Antalya, Türkiye kıyılarından yakalanan *Lagocephalus sceleratus* (Gmelin, 1789)'in MS/MS ve GC/MS analizleriyle tetrodotoksin ve yağ asitleri içeriği

Öz

Türkiye kıyılarından Antalya'da yakalanan beş *Lagocephalus sceleratus* (Gmelin, 1789)'in MS/MS ve GC/MS analizleriyle tetrodotoksin ve yağ asitleri içeriği çalışıldı. LC-MS/MS analizi ile barsak, karaciğer, ovar ve kasta TTX bulundu. GC/MS analizleriyle yağ asitleri tespit edildi. Yağ içeriği yüksek değil fakat çoklu doymamış n-3 and n-6 oranları yüksektir.

Anahtar Kelimeler: TTX, tetrodotoksin, yağ asitleri, *Lagocephalus sceleratus*, LC-MS/MS, GC/MS.

References

- Adams, R.P. (2007) Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry. Vol., Allured Publ. Corp, Carol Stream, IL.
- Akyol, O., Ünal, V., Ceyhan, T., Bilecenoğlu, M. (2005) First confirmed record of *Lagocephalus sceleratus* (Gmelin, 1789) in the Mediterranean Sea. *J. Fish Biol.* 66: 1183-1186.
- Aslan, S.S., Coban, B., Tekinay, A., Gezgin, T., Güven, K.C. (2009) Effects of pellet and extruded feed on fatty acid composition of European Sea Bass (*Dicentrarchus labrax*) using HRGC-MS. *Fres. Environ. Bull.* 18: 112-116.

- Aydin, M. (2011) Growth, reproduction and diet of pufferfish (*Lagocephalus sceleratus* Gmelin, 1789) from Turkey's Mediterranean Sea coast. *Turk. J. Fish. Aquat. Sci.* 11: 589-596.
- Aydin, M., Tufan, B., Sevgili, H., Köse, S. (2013) Seasonal changes in proximate composition and fatty acid profile of Pufferfish (*Lagocephalus sceleratus* Gmelin, 1789) from the Mediterranean Sea of Turkey. *J. Aquat. Food Prod T.* 22: 178-191.
- Bane, V., Lehane, M., Dikshit, M., Riordan, A., Furey, A. (2014) Tetrodotoxin: chemistry, toxicity, source, distribution and detection. *Toxins* 6: 693.
- Bentur, Y., Ashkar, J., Lurie, Y., Levy, Y., Azzam, Z.S., Litmanovich, M., Golik, M., Gurevych, B., Golani, D., Eisenman, A. (2008) Lessepsian migration and tetrodotoxin poisoning due to *Lagocephalus sceleratus* in the eastern Mediterranean. *Toxicon.* 52: 964-968.
- Brillantes, S., Samosorn, W., Faknoi, S., Oshima, Y. (2003) Toxicity of puffers landed and marketed in Thailand. *Fisheries Sci.* 69: 1224-1230.
- Chen, C.Y., Chou, H.N. (1998) Detection of tetrodotoxin by high performance liquid chromatography in lined-moon shell and puffer fish. *Acta Zoologica Taiwanica* 9: 41-48.
- Chulanetra, M., Sookrung, N., Srimanote, P., Indrawattana, N., Thanongsaksrikul, J., Sakolvaree, Y., Chongsa-Nguan, M., Kurazono, H., Chaicumpa, W. (2011) Toxic marine puffer fish in Thailand Seas and tetrodotoxin they contained. *Toxins* 3: 1249-1262.
- Curvers, J., Rijks, J., Cramers, C., Knauss, K., Larson, P. (1985) Temperature programmed retention indices: Calculation from isothermal data. Part 1: Theory. *J. High Res. Chromatog.* 8: 607-610.
- Daviglus, M.L., Stamler, J., Orenca, A.J., Dyer, A.R., Liu, K., Greenland, P., Walsh, M.K., Morris, D., Shekelle, R.B. (1997) Fish consumption and the 30-year risk of fatal myocardial infarction. *New Engl. J. Med.* 336: 1046-1053.
- Golani, D. (2002) Lessepsian Fish Migration, Characterization and Impact on the Eastern Mediterranean. In: Workshop on Lessepsian Migration, 20–21 July 2002 (eds., B. Öztürk, N. Başusta) Gökçeada, Turkey, pp.1-9.
- Goto, T., Kishi, Y., Takahashi, S., Hirata, Y. (1965) Tetrodotoxin. *Tetrahedron* 21: 2059-2088.

- Hochmuth, H. (2008) MassFinder-4. Vol., Hochmuth Scientific Consulting, Hamburg, Germany.
- Hwang, P.A., Tsai, Y.H., Lin, S.J., Hwang, D.F. (2007) The gastropods possessing TTX and/or PSP. *Food Rev. Int.* 23: 321-340.
- Jang, J.H., Lee, J.S., Yotsu-Yamashita, M. (2010) LC/MS Analysis of tetrodotoxin and its deoxy analogs in the marine puffer fish fugu niphobles from the southern coast of Korea, and in the brackishwater puffer fishes tetraodon nigroviridis and tetraodon biocellatus from Southeast Asia. *Mar. Drugs* 8: 1049.
- Jen, H.C., Lin, S.J., Tsai, Y.H., Chen, C.H., Lin, Z.C., Hwang, D.F. (2008) Tetrodotoxin poisoning evidenced by solid-phase extraction combining with liquid chromatography-tandem mass spectrometry. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* 871: 95-100.
- Joseph, J.D., Ackman, R.G. (1992) Capillary column gas chromatography method for analysis of encapsulated fish oil and fish oil ethyl esters: collaborative study. *J. AOAC Int.* 75: 488-506.
- Kızılkaya, Z. (2014) Why is it important monitoring climate change in MPAs? Monitoring of *Lagocephalus sceleratus* in Turkish MPAs/Case studies. In: The Network of Marine Protected Areas Managers in Mediterranean, Climate Change Training Workshop, Cadaques, Spain.
- Kono, M., Matsui, T., Furukawa, K., Yotsu-Yamashita, M., Yamamori, K. (2008) Accumulation of tetrodotoxin and 4,9-anhydrotetrodotoxin in cultured juvenile kusafugu *Fugu niphobles* by dietary administration of natural toxic komonfugu *Fugu poecilonotus* liver. *Toxicon.* 51: 1269-1273.
- Köşker, R. (2014) Captured in Mersin Bay blowfish (*Lagocephalus sceleratus*) and the gender depending on season, nutrient composition and tetrodotoxin (TTX) in the level of the level of analysis. Institute of Science, Çukurova University, Adana.
- Köşker, A.R., Özoğul, F., Ayaş, D., Durmuş, D., Uçar, Y. (2015) The new toxin of Mediterranean:Tetrodotoxin. *Ege J. Fish. Aqua. Sci.* 32: 15-24.
- Lin, W., Hwang, D. (2012) Analysis of poisoning cases, monitoring and risk warning for marine toxins (TTX, PSP and CTXs) in Taiwan. *J. Food Drug Anal.* 20: 764-771.
- Love, R.M. (1970) The Chemical Biology of Fishes. Vol., Academic Press, London & New York.

McLafferty, F.W., Stauffer, D.B. (1989) The Wiley/NBS Registry of Mass Spectral Data. Vol., J. Wiley and Sons, New York.

Miyazawa, K., Noguchi, T. (2001) Distribution and origin of tetrodotoxin. *J. Toxicol. Toxin Rev.* 20: 11-33.

Nader, M.R., Indary, S., Boustany, L.E. (2012) The puffer fish *Lagocephalus sceleratus* (Gmelin, 1789) in the eastern Mediterranean. In: EastMed Technical Documents 10 (ed., FAO), FAO UN, Athens, Greece.

Noguchi, T., Arakawa, O. (2008) Tetrodotoxin - distribution and accumulation in aquatic organisms, and cases of human intoxication. *Mar. Drugs* 6: 220-242.

Nurullohoğlu, Z.U., Ulusoy, E. (2011) Fatty acid composition of *Lagocephalus sceleratus* (Gmelin, 1789) (Osteichthyes: Tetraodontidae). *Marmara University Journal of science* 25: 184-191.

Oyaizu, M., Fujimoto, Y., Takenaga, F., Itoh, S. (2000) Fatty acid composition of total lipids in puffer fish meat. *Food Preserv. Sci.* 26: 333-338.

Özoğul, Y., Özoğul, F.H., Çiçek, E., Polat, A., Kuley, E. (2009) Fat content and fatty acid compositions of 34 marine water fish species from the Mediterranean Sea. *Int. J. Food Sci. Nutr.* 60: 464-475.

Rodriguez, P., Alfonso, A., Otero, P., Katikou, P., Georgantelis, D., Botana, L.M. (2012) Liquid chromatography–mass spectrometry method to detect tetrodotoxin and its analogues in the puffer fish *Lagocephalus sceleratus* (Gmelin, 1789) from European waters. *Food Chem.* 132: 1103-1111.

Sabrah, M.M., El-Ganainy, A.A., Zaky, M.A. (2006) Biology and toxicity of the puffer fish *Lagocephalus sceleratus* (Gmelin, 1789) from the Gulf of Suez. *Egypt. J. Aquatic Res.* 32: 283-297.

Saoudi, M., Abdelmouleh, A., Kammoun, W., Ellouze, F., Jamoussi, K., El Feki, A. (2008) Toxicity assessment of the puffer fish *Lagocephalus lagocephalus* from the Tunisian coast. *C. R. Biol.* 331: 611-616.

Shoji, Y., Yotsu-Yamashita, M., Miyazawa, T., Yasumoto, T. (2001) Electrospray Ionization Mass Spectrometry of Tetrodotoxin and Its Analogs: Liquid Chromatography/Mass Spectrometry, Tandem Mass Spectrometry, and Liquid Chromatography/Tandem Mass Spectrometry. *Anal. Biochem.* 290: 10-17.

Silva, M., Azevedo, J., Rodriguez, P., Alfonso, A., Botana, L.M., Vasconcelos, V. (2012) New gastropod vectors and tetrodotoxin potential expansion in temperate waters of the Atlantic Ocean. *Mar. Drugs* 10: 712-726.

Simidu, U., Kitatsukamoto, K., Yasumoto, T. (1990) Taxonomy of four marine bacterial strains that produce tetrodotoxin. *Int. J. Syst. Bacteriol.* 40: 331-336.

Simon, K.D., Mazlan, A.G., Usup, G. (2009) Toxicity of puffer fishes (*Lagocephalus wheeleri* Abe, Tabeta and Kitahama, 1984 and *Lagocephalus sceleratus* Gmelin, 1789) from the East coast waters of Peninsular Malaysia. *J. Biol. Sci.* 9: 482-487.

Tsuda, K., Ikuma, S., Kawamura, M., Tachikawa, R., Sakai, K., Tamura, C., Amakasu, D. (1964) Tetrodotoxin. VII. On the structures of tetrodotoxin and its derivatives. *Chem. Pharm. Bull.* 12: 1357-1374.

Woodward, R.B. (1964) The structure of tetrodotoxin. *Pure Appl. Chem.* 9: 49-74.

Wu, Z., Xie, L., Xia, G., Zhang, J., Nie, Y., Hu, J., Wang, S., Zhang, R. (2005) A new tetrodotoxin-producing actinomycete, *Nocardiopsis dassonvillei*, isolated from the ovaries of puffer fish *Fugu rubripes*. *Toxicon.* 45: 851-859.

Yokoo, A. (1950) Chemical studies on pufferfish toxin (3) - separation of spheroidine. *Nippon Kagaku Zasshi* 71: 590-592.

Yu, V.C.H., Yu, P.H.F., Ho, K.C., Lee, F.W.F. (2011) Isolation and identification of a new tetrodotoxin-producing bacterial species, *Raoultella terrigena*, from Hong Kong marine puffer fish *Takifugu niphobles*. *Mar. Drugs* 9: 2384.